



MASTER GARDENER

COLORADO STATE UNIVERSITY
EXTENSION

CMG GardenNotes #136

Plant Structures: Fruit

Outline: Function, page 1
 Structure, page 1
 Fruit types, page 2
 Fruit growth terms, page 3



Thought question:

- o Why are fading flowers removed from spring flowering bulbs and other flowering ornamental plants?
-

Fruit evolves from the maturing ovary following pollination and fertilization. Fruits can be either fleshy or dry. They contain one or more seeds.

Function

- Reproduction
- Horticulture uses
 - Feed, food, and oils
 - Aesthetic qualities
 - Plant identification

Structure

Fruit consists of carpels where the ovules (seeds) develop and the ovary wall or **pericarp**, which may be fleshy (as in apples) or dry and hard (as in an acorn). Some fruits have seeds (mature ovules) enclosed within the ovary (apples, peaches, oranges, squash and cucumbers). The peel of an orange, the pea pod, the sunflower shell, and the skin flesh and pit of a peach are all derived from the pericarp.

Other fruit have seeds that are situated on the periphery of the pericarp (corn cob, strawberry flesh).

Figure 1. In apples, the ovary wall becomes the fleshy part of the fruit. Notice the small fruit structure in the blossom.






Figure 2. Pome fruit (apple)

Figure 3. Stone fruit (peach)




Fruit Types

Conifers

Conifers are best known for their woody cones, pinecones. Junipers are an example of a conifer with a fleshy cone (juniper berry). Upon close examination, the overlapping scales can be observed.

Figure 4. Fruit of conifers – Left: Woody seed cone (pinecone). Right: Fleshy seed cone (Juniper berry).



Flowering Plants

Depending on flower structure and inflorescence type, fruits may be either simple, aggregate, or multiple.

Accessory- fruit having some flesh derived from tissue exterior to the carpel.

Simple – Fruit formed from one ovary.

Aggregate – Fruit formed from a single flower with many ovaries. If not all ovaries are pollinated and fertilized, fruit will be misshapen (raspberry, magnolia).

Multiple – Fruit developed from a fusion of separate, independent flowers borne on a single structure (mulberry, pineapple, beet seed).

Fruit Growth Terms

Pollination – Transfer of pollen from the male flower to the stigma of the female flower.

Fertilization – Union of the pollen grain from the male flower with the egg cell in the female flower.

Drop – Fruit drops when not pollinated or fertilized and when too much fruit sets on a tree.

Growth – Primarily cell enlargement as the cells fill with water.

Climacteric – Point when a fruit will continue to ripen if removed from a plant; for example, pumpkins turning orange after being harvested.

Authors: David Whiting, Consumer Horticulture Specialist (retired), Colorado State University Extension; with Michael Roll and Larry Vickerman (former CSU Extension employees). Line drawings by Scott Johnson and David Whiting. Revised by Patti O'Neal, Roberta Tolan and Mary Small, CSU Extension.

- Colorado Master Gardener *GardenNotes* are available online at www.cmg.colostate.edu.
- Colorado Master Gardener training is made possible, in part, by a grant from the *Colorado Garden Show, Inc.*
- Colorado State University, U.S. Department of Agriculture and Colorado counties cooperating.
- Extension programs are available to all without discrimination.
- No endorsement of products mentioned is intended nor is criticism implied of products not mentioned.
- Copyright Colorado State University Extension. All Rights Reserved. *CMG GardenNotes* may be reproduced, without change or additions, for nonprofit educational use with attribution.

Revised July 2016